

DATASHEET

Scedosporium, mouse monoclonal antibody [clone HG12]

Product code	ID2510-0025 / ID2510-0100
Unit size	25µL / 100µL
Concentration	See batch specific Certificate of Analysis
Alternative name	No
Clone	HG12
Host	Mouse
Antigen	120kDa glycoprotein
Isotype	lgG1
Purification	Protein A
Applications	ELISA, DAS-ELISA, WB, IF, IEM For DAS-ELISA, use in combination with mAb GA3 [ID2515]
Recommended dilution	1:1000
Optimisation	Optimal dilutions to be determined by end user
Species reactivity	Pseudallescheria boydii, Scedosporium boydii, Scedosporium apiospermum, Scedosporium aurantiacum
Storage buffer	Phosphate buffered saline pH7.2 with 0.095% (w/v) sodium azide
Shipping	Blue ice
Storage temperature	Store as supplied at $+2^{\circ}C \sim +8^{\circ}C$ for up to 1 year
Reference	Thornton CR. (2009). Tracking the emerging human pathogen <i>Pseudallescheria boydii</i> by using highly specific monoclonal



Legend: Immunogold electron microscopy (IEM) showing distribution of HG12 antigen in cell wall and cytoplasm of *P. boydii* spore following phagocytosis by a mouse alveolar macrophage. Tissue section was probed with mAb HG12 followed by goat anti-mouse 20nm gold conjugate.

antibodies. Clinical and Vaccine Immunology 16: 756-764.

UNLESS OTHERWISE STATED IN OUR CATALOG OR OTHER COMPANY DOCUMENTATION ACCOMPANYING THE PRODUCT(S), OUR PRODUCTS ARE INTENDED FOR **RESEARCH USE** ONLY (RUO) AND ARE NOT TO BE USED FOR ANY OTHER PURPOSE, WHICH INCLUDES, BUT IS NOT LIMITED TO, UNAUTHORISED COMMERCIAL USES, *IN VITRO* DIAGNOSTIC USES, *EX* VIVO OR *IN VIVO* THERAPEUTIC USES OR ANY TYPE OF CONSUMPTION OR APPLICATION TO HUMANS OR ANIMALS.

> ISCA Diagnostics Ltd., Floor 3, Geoffrey Pope Building, Stocker Road, Exeter, EX4 4QD. techsupport@iscadiagnostics.com | Phone: +44/0 7527 869 861